



DOCUMENTATION OF HYPOGLYCEMIA TREATMENT PROTOCOL


(file this form within the Progress Note Section of the Chart)

IMPRINT PATIENT IDENTIFICATION HERE

General Information:

- Check fingerstick BG if patient experiences any symptoms of hypoglycemia including Pallor, Clammy Skin, Hunger, Restless Sleep, Fatigue, Headache, Confusion, Sweatiness and Irritability
- If bedside BG <70, confirm by repeating bedside test immediately as per policy, and **Do Not Delay Hypoglycemia Treatment**
- Check all IV infusions to ensure all medications (including IV insulin, if applicable) are being administered as ordered.
- If the patient is/was taking oral medication for diabetes in the sulfonylurea class (glipizide [Glucotrol®], glyburide [Glynase®, Micronase®, DiaBeta®]; glimepiride [Amaryl®], chlorpropamide [Diabinese®], tolazamide [Tolinase®]) **then hold the sulfonylurea, follow this algorithm, check BG every 4 hours X 24 hours and consider immediate Diabetes Consult (Pager #1082)**
- **If this is the second BG <40 in the past 24-48 hours, notify attending service**

Document low Blood Glucose (BG): _____ Date: _____ Time: _____ RN: _____



* If glucagon is used, it should be followed as soon as possible by EITHER oral carbohydrates (if patient can subsequently eat) or by obtaining IV access and starting D5W as indicated in the box. Glucagon should not be repeated.

